

# IOWA STATE UNIVERSITY

## Digital Repository

---

Chemistry Publications

Chemistry

---

5-16-2018

## The first synthesis of biatriosporin D

Shuai Wang

*Iowa State University*, [swang24@iastate.edu](mailto:swang24@iastate.edu)

George A. Kraus

*Iowa State University*, [gakraus@iastate.edu](mailto:gakraus@iastate.edu)

Follow this and additional works at: [https://lib.dr.iastate.edu/chem\\_pubs](https://lib.dr.iastate.edu/chem_pubs)



Part of the [Organic Chemistry Commons](#)

The complete bibliographic information for this item can be found at [https://lib.dr.iastate.edu/chem\\_pubs/1075](https://lib.dr.iastate.edu/chem_pubs/1075). For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

---

This Article is brought to you for free and open access by the Chemistry at Iowa State University Digital Repository. It has been accepted for inclusion in Chemistry Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact [digirep@iastate.edu](mailto:digirep@iastate.edu).

---

# The first synthesis of biatriosporin D

## Abstract

Biatriosporin D was synthesized in six steps from 3,5-dimethoxyphenol. The key step was an intramolecular Friedel-Crafts cyclization of an acyl imidazolide to form an ortho-quinone.

## Keywords

Biatriosporin D, Friedel-Crafts cyclization, Ortho-quinone

## Disciplines

Organic Chemistry

## Comments

This is a manuscript of an article published as Wang, Shuai, and George A. Kraus. "The first synthesis of biatriosporin D." *Tetrahedron Letters* 59, no. 20 (2018): 1968-1969. DOI: [10.1016/j.tetlet.2018.04.027](https://doi.org/10.1016/j.tetlet.2018.04.027).  
Posted with permission.

## Creative Commons License



This work is licensed under a [Creative Commons Attribution-Noncommercial-No Derivative Works 4.0 License](https://creativecommons.org/licenses/by-nc-nd/4.0/).


## Graphical Abstract

To create your abstract, type over the instructions in the template box below.  
Fonts or abstract dimensions should not be changed or altered.

### The First Synthesis of Biatriosporin D

Shuai Wang, George A. Kraus

Leave this area blank for abstract info.



overall yield 50 % in 6 steps

# The First Synthesis of Biatriosporin D

Shuai Wang <sup>a</sup>, George A. Kraus <sup>a, \*</sup>

<sup>a</sup> Department of Chemistry, Iowa State University, Ames, IA 50010, U.S.A.

## ARTICLE INFO

### Article history:

Received

Received in revised form

Accepted

Available online

### Keywords:

Biatriosporin D

Friedel-Crafts cyclization

Ortho-quinone


## ABSTRACT

Biatriosporin D was synthesized in six steps from 3,5-dimethoxyphenol. The key step was an intramolecular Friedel-Crafts cyclization of an acyl imidazolide to form an ortho-quinone.

The biatriosporins are a group of heptaketide metabolites isolated from the endolichenic fungus *Biatrospora* sp.<sup>1</sup> They have been reported to exhibit antifungal activity. Recently, Lou and coworkers reported that biatriosporin D (**1**) shows anti-virulence activity by a mechanism leading to the reduction of intracellular cAMP levels.<sup>2</sup> In the context of our program in antimicrobials,<sup>3</sup> we needed a sample of biatriosporin D and herein report a direct synthetic route to **1**.

resulted in two products in approximately equal yields. The structure of compound **7** was supported by the strong NOE between the furan hydrogen and the methyl group on the furan ring. The structure of compound **6** was supported by the symmetry observed in both the proton NMR spectra. Although compound **7** could have been converted into **1**, the yield of **7** was low. Therefore, this strategy was abandoned in favor of the plan


**Figure 1.** Biatriosporins




approach focused on creating the ortho-quinone subunit in **1** via a Friedel-Crafts reaction with oxalyl chloride in Scheme 1. We obtained the requisite aryl furan from the palladium mediated coupling of the triflate of 3,5-dimethoxyphenol (**3**) with 5-methylfuran. Unfortunately, the key acylation with oxalyl chloride was not selective, providing several decomposition products. Various Lewis acids (BF<sub>3</sub>, AlCl<sub>3</sub>, TiCl<sub>4</sub>, SnCl<sub>4</sub>) were tried but none of them afforded the desired product in good yield. Despite changes in reaction parameters such as solvent (CH<sub>2</sub>Cl<sub>2</sub>, THF, 1,4-dioxane), temperature and the addition of reagents to capture HCl (propylene oxide) the best yield was only 10% with titanium tetrachloride in methylene chloride at -78 °C.

### Scheme 1. Friedel-Crafts reaction with oxalyl chloride.

Next, we explored the reaction between **4** and ethyl chlorooxoacetate catalyzed by AlCl<sub>3</sub> in Scheme 2.<sup>4</sup> The reaction



**Scheme 2.** Friedel-Crafts reaction with ethyl chlorooxoacetate.



\* Corresponding author. Tel.: +1-515-294-7794; e-mail: gakraus@iastate.edu

## A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at

## References and notes

1. Zhou, Y.; Zhang, M.; Zhu, R.X.; Zhang, J.Z.; Xie, F.; Li, X.B.; Chang, W.Q.; Wang, X.N.; Zhao, Z.T.; Lou, H.X. *J. Nat. Prod.* 2016, 79, 2149-2157.
2. Zhang, M.; Chang, W.Q.; Shi, H.Z.; Zhou, Y.H.; Zheng, S.; Li, Y.; Li, L.; Lou, H.X. *Toxicology and Applied Pharmacology*. 2017, 322, 104-112.
3. Guney, T.; Kohles, S.A.; Thompson, V.L.; Philips, G.J.; Kraus, G.A. *Tetrahedron*. 2015, 71, 3115-3119.
4. Richter, C.; Ernsting, N.P.; Mahrwald, R. *Synthesis*. 2016, 48, 1217-1225.
5. Kraus, G.A.; Gupta, V.; Kempenma, A. *Synlett*. 2012, 23, 385-388.
6. Yoshikawa, N.; Doyle, A.; Tan, L.; Murry, J.A.; Akao, A.; Kawasaki, M.; Sato, K. *Org. Lett.* 2007, 9, 4103-4106.
7. Bernard, A.M.; Ghiani, M.R.; Piras, P.P.; Rivoldini, A. *Synthesis*. 1989, 287-289.
8. **General Experimental Procedure for the cyclization and the demethylation.** To a solution of acid **10** (145 mg, 1.0 equiv) in CH<sub>2</sub>Cl<sub>2</sub> (3.0 mL) was added carbonyldiimidazole (162 mg, 2.0 equiv). The reaction mixture was stirred at room temperature overnight. The resulting solution was added into a solution of TiCl<sub>4</sub> (380 mg, 4.0 equiv) in CH<sub>2</sub>Cl<sub>2</sub> (2.0 mL) at -10 °C. The reaction mixture was stirred at -10 °C for 5 hours and quenched by addition of HCl (2.0 M) and then was extracted twice with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic solution was washed with sat NaHCO<sub>3</sub>, water and brine. The organic solution was dried by Na<sub>2</sub>SO<sub>4</sub>. The solvent was evaporated in vacuo and red solid **5** was obtained (87 mg, 63 %). To a solution of solid **5** (54.4 mg, 1.0 equiv) in DMF (6.0 mL) was added LiCl (50.4 mg, 6.0 equiv). The reaction mixture was heated under reflux for 2 hours and was cooled down. DMF was removed by distillation under reduced pressure. And the residue was acidified by HCl (2.0 M) and then was extracted three times by EtOAc. The combined organic solution was washed with water, brine. The organic solution was dried by Na<sub>2</sub>SO<sub>4</sub>. The solvent was evaporated in vacuo and the residue was purified by column chromatography on silica gel and product **1** was obtained as red solid (47.4 mg, 92 %). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ = 12.43 (s, 1H), 6.74 (d, J=2.3, 1H), 6.41 (s, 1H), 6.31 (d, J=2.3, 1H), 3.90 (s, 3H), 2.40 (s, 3H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ = 182.12, 175.37, 169.22, 168.20, 158.14, 156.31, 129.59, 122.78, 106.82, 105.39, 104.44, 101.26, 56.31, 13.78. HRMS (ESI-QTOF) calcd for [M + H<sup>+</sup>]: 259.0601, found: 259.0597.

Scheme 3. Synthesis of **1** in six steps.

Our second route began with the reaction of 3,5-dimethoxyphenol with ethyl chlorooxalate and titanium tetrachloride to produce the phenol,<sup>5</sup> which could be readily converted into triflate **8** in 93 % yield. Coupling of **8** with 5-methylfuran boronic acid afforded furan **9** in 94 % isolated yield. Hydrolysis with NaOH in ethanol-water provided the keto acid **10** in 99% yield. Conversion of the keto acid **10** into ortho-quinone **5** proved to be difficult. Reaction with trifluoroacetic anhydride and tin tetrachloride produced 12-20% yields of **5**. Fortunately, a procedure recently reported by Yoshikawa<sup>6</sup> using carbonyl diimidazole (CDI) followed by titanium tetrachloride at -10 °C afforded a 63% isolated yield of **5** as a sparingly soluble red solid. Attempted deprotection using boron trichloride, boron tribromide or aluminum chloride returned recovered starting material. Finally, lithium chloride in boiling DMF for two hours produced the demethylation product **1** in 92 % yield.<sup>7</sup>

Ortho-quinone **1** was synthesized in six steps from 3,5-dimethoxyphenol. This route is scalable, providing 300 milligrams of **1**. This three-component synthetic approach will allow for considerable flexibility in the design of synthetic analogs of **1**.

## Acknowledgments

We thank the Department of Chemistry for partial support of Shuai Wang.